



Anabolic Steroid E Testosterone Powder For Bodybuilding CAS 315-37-

7

Our Product Introduction

for more products please visit us on peptide-powder.com


Basic Information

- Place of Origin: China
- Brand Name: Hongbaiyi
- Certification: COA, HPLC MR
- Model Number: HBY-Testosterone
- Minimum Order Quantity: 1KG
- Price: Negotiable
- Packaging Details: Aluminum Bag or 25KG/Febr Drum
- Delivery Time: 3-5 work days after your payment
- Payment Terms: , Western Union, T/T
- Supply Ability: 4000kg/Month



Product Specification

- Product: Testosterone Enanthate
- MF: C₂₆H₄₀O₃
- MW: 400.6
- Cas No.: 315-37-7
- Apperance: White Crystalline Powder
- Purity: 99%Min
- Storage Temp: 2-8°C
- Highlight: **Anabolic E Testosterone Powder,**
315-37-7 E Testosterone Powder,
315-37-7 test ethanate cycle



More Images



Product Description

Testosterone Anabolic Steroid Testosterone Enanthate For Bodybuilding CAS 315-37-7

Description Of Testosterone Enanthate

Testosterone Enanthate Injection, USP provides Testosterone Enanthate, USP, a derivative of the primary endogenous androgen testosterone, for intramuscular administration. In their active form, androgens have a 17-beta-hydroxy group. Esterification of the 17-beta-hydroxy group increases the duration of action of testosterone; hydrolysis to free testosterone occurs in vivo. Each mL of sterile, colorless to pale yellow, solution provides 200 mg Testosterone Enanthate, USP in sesame oil with 5 mg chlorobutanol (chloral derivative) as a preservative.

Testosterone Enanthate, USP is designated chemically as androst-4-en-3-one, 17-[(1-oxoheptyl)-oxy]-, (17 β)-.

Testosterone Enanthate - Clinical Pharmacology

Endogenous androgens are responsible for the normal growth and development of the male sex organs and for maintenance of secondary sex characteristics. These effects include growth and maturation of prostate, seminal vesicles, penis, and scrotum; development of male hair distribution, such as beard, pubic, chest, and axillary hair; laryngeal enlargement; vocal chord thickening; alterations in body musculature; and fat distribution.

Androgens also cause retention of nitrogen, sodium, potassium, and phosphorus, and decreased urinary excretion of calcium. Androgens have been reported to increase protein anabolism and decrease protein catabolism. Nitrogen balance is improved only when there is sufficient intake of calories and protein.

Androgens are responsible for the growth spurt of adolescence and for the eventual termination of linear growth which is brought about by fusion of the epiphyseal growth centers. In children, exogenous androgens accelerate linear growth rates but may cause a disproportionate advancement in bone maturation. Use over long periods may result in fusion of the epiphyseal growth centers and termination of the growth process. Androgens have been reported to stimulate the production of red blood cells by enhancing the production of erythropoietic stimulating factor.

During exogenous administration of androgens, endogenous testosterone release is inhibited through feedback inhibition of pituitary luteinizing hormone (LH). At large doses of exogenous androgens, spermatogenesis may also be suppressed through feedback inhibition of pituitary follicle stimulating hormone (FSH).

Testosterone enanthate is used for

Treating symptoms of low testosterone in men when the body does not make any testosterone or not enough testosterone (hypogonadism). It may also be used for other conditions as determined by your doctor. It is also used to treat certain types of breast cancer in women.

Testosterone enanthate is a male sex hormone. It works by replacing or supplementing the testosterone that is naturally made in the body. It also counteracts the effects of estrogen in certain types of breast cancer, which helps to decrease cancer growth.

Basic Information form Of Testosterone Enanthate:

Product Name	Testosterone Enanthate
Cas No.	315-37-7
Purity	99%min
Appearance	White Crystalline powder
MF	C ₂₆ H ₄₀ O ₃
MW	400.6
Storage Temp	2-8°C
Melting point	34 ~ 39°

Product Image Of Testosterone Enanthate



Do NOT use testosterone enanthate if

you are allergic to any ingredient in testosterone enanthate

you are pregnant, may become pregnant, or are breast-feeding

you have breast cancer and are male; known or suspected prostate cancer; or serious heart, liver, or kidney problems

Contact your doctor or health care provider right away if any of these apply to you.

Before using testosterone enanthate

Some medical conditions may interact with testosterone enanthate. Tell your doctor or pharmacist if you have any medical conditions, especially if any of the following apply to you:

if you are pregnant, planning to become pregnant, or are breast-feeding

if you are taking any prescription or nonprescription medicine, herbal preparation, or dietary supplement

if you have allergies to medicines, foods, or other substances (especially sesame oil)

if you have heart disease, heart failure, coronary artery disease, angina (chest pain), high cholesterol levels, swelling (edema), lung disease, or sleep apnea (long pauses in breathing while you sleep)

if you have diabetes, an enlarged prostate, kidney or liver disease, high blood calcium levels, or obesity

if you have a history of blood clots

Some MEDICINES MAY INTERACT with testosterone enanthate. Tell your health care provider if you are taking any other medicines, especially any of the following:

Carbamazepine, corticosteroids (eg, prednisone), macrolide immunosuppressants (eg, tacrolimus), or oxyphenbutazone because their actions and the risk of their side effects may be increased by testosterone enanthate

Anticoagulants (eg, warfarin), insulin, or oral hypoglycemics (eg, glyburide) because their side effects, including risk of bleeding, may be increased by testosterone enanthate

Propranolol because its effectiveness may be decreased by testosterone enanthate

This may not be a complete list of all interactions that may occur. Ask your health care provider if testosterone enanthate may interact with other medicines that you take. Check with your health care provider before you start, stop, or change the dose of any medicine.

How to use testosterone enanthate

Use testosterone enanthate as directed by your doctor. Check the label on the medicine for exact dosing instructions.

Testosterone enanthate is usually given as an injection at your doctor's office, hospital, or clinic. If you will be using testosterone enanthate at home, a health care provider will teach you how to use it. Be sure you understand how to use testosterone enanthate. Follow the procedures you are taught when you use a dose. Contact your health care provider if you have any questions.

Do not use testosterone enanthate if it contains particles, is cloudy or discolored, or if the vial is cracked or damaged.

Keep this product, as well as syringes and needles, out of the reach of children and away from pets. Do not reuse needles, syringes, or other materials. Ask your health care provider how to dispose of these materials after use. Follow all local rules for disposal.

If you miss a dose of testosterone enanthate, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

Ask your health care provider any questions you may have about how to use testosterone enanthate.

Testosterone Enanthate Dosage and Administration

Prior to initiating Testosterone Enanthate injection, confirm the diagnosis of hypogonadism by ensuring that serum testosterone concentrations have been measured in the morning on at least two separate days and that these serum testosterone concentrations are below the normal range.

Dosage and duration of therapy with Testosterone Enanthate injection will depend on age, sex, diagnosis, patient's response to treatment, and appearance of adverse effects. When properly given, injections of Testosterone Enanthate, are well tolerated. Care should be taken to slowly inject the preparation deeply into the gluteal muscle, being sure to follow the usual precautions for intramuscular administration, such as the avoidance of intravascular injection (see PRECAUTIONS).

In general, total doses above 400 mg per month are not required because of the prolonged action of the preparation. Injections more frequently than every two weeks are rarely indicated. NOTE: Use of a wet needle or wet syringe may cause the solution to become cloudy; however this does not affect the potency of the material. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Testosterone Enanthate injection is a clear, colorless to pale yellow solution.

Male hypogonadism: As replacement therapy, i.e., for eunuchism, the suggested dosage is 50 to 400 mg every 2 to 4 weeks.

In males with delayed puberty: Various dosage regimens have been used; some call for lower dosages initially with gradual increases as puberty progresses, with or without a decrease to maintenance levels. Other regimens call for higher dosage to induce pubertal changes and lower dosage for maintenance after puberty. The chronological and skeletal ages must be taken into consideration, both in determining the initial dose and in adjusting the dose. Dosage is within the range of 50 to 200 mg every 2 to 4 weeks for a limited duration, for example, 4 to 6 months. X-rays should be taken at appropriate intervals to determine the amount of bone maturation and skeletal development (see INDICATIONS AND USAGE and WARNINGS).

Palliation of inoperable mammary cancer in women: A dosage of 200 to 400 mg every 2 to 4 weeks is recommended. Women with metastatic breast carcinoma must be followed closely because androgen therapy occasionally appears to accelerate the disease.

Package and Shipping Of Testosterone Enanthate:

Under 25kg: inner: A PE bag in aluminum foil bag; Outside: carton

Over 25kg: inner: Double PE bags; Outside: 25kg/ fibre drum

Shipping:

With 3-5 days after your payment



FAQ Of Testosterone Enanthate:

Q1: Do you Accept Sample Order?

A: Yes. The smallest order we accept is 1kg.

Q2: Do you accept VISA business credit card?

A: Yes. VISA business credit card is in our payment.

Q3: Is there any discount?

A: Yes, if you want to purchase larger quantity, we can offer a better price for you.

Q4: How long does it take to me if I order?

A: About 3-5 days upon receipt of payment.

Q5: What kind of transportation do you have?

A: We have many ways of transportations, eg: Air, Sea, Special Lines, TNT, FEDEX, EMS, ect.



Shaanxi Hongbaiyi Biotech Co., Ltd.



18192109180



tracy@sxhongbaiyi.com



peptide-powder.com

