



Sildenafil Citrate 100mg Sex Drug Male Enhancement Steroids Revatio CAS 171599-83-0

Our Product Introduction

for more products please visit us on peptide-powder.com

Basic Information

- Place of Origin: China
- Brand Name: Hongbaiyi
- Certification: COA, HPLC MR
- Model Number: HBY-Sildenafil
- Minimum Order Quantity: 1KG
- Price: Negotiable
- Packaging Details: Aluminum Bag or 25KG/Febr Drum
- Delivery Time: 3-5 work days after your payment
- Payment Terms: , Western Union, T/T
- Supply Ability: 4000kg/Month



Product Specification

- Product Name: Sildenafil Citrate 100mg
- Cas No: 171599-83-0
- Synonyms: Sildenafil Citrate Sildenafil
- MF: C₂₈H₃₈N₆O₁₁S
- MW: 666.7
- Purity: >99%
- Appearance: White Crystals Powder
- Shelf Life: 2 Years
- Highlight: Sildenafil Citrate 100mg Sex Drug, Steroids Revatio Male Enhancement Powder, CAS 171599-83-0 Sildenafil Citrate 100mg



More Images



Product Description

Sildenafil Citrate 100mg Sex Drug Male Enhancement Steroids Revatio CAS 171599-83-0



Basic Information form Of Sildenafil Citrate 100mg:

Name	Sildenafil Citrate 100mg
Cas No.	171599-83-0
Synonyms	Sildenafil Citrate,Sildenafil
MF	C28H38N6O11S
MW	666.7
Appearance	white crystals powder
Purity	>99%
Shelf life	2 years

What Is Sildenafil Citrate 100mg?

Sildenafil is used to treat erectile dysfunction (impotence; inability to get or keep an erection) in men. Sildenafil (Revatio) is used to improve the ability to exercise in adults with pulmonary arterial hypertension (PAH; high blood pressure in the vessels carrying blood to the lungs, causing shortness of breath, dizziness, and tiredness). Children should not usually take sildenafil, but in some cases, a doctor may decide that sildenafil (Revatio) is the best medication to treat a child's condition. Sildenafil is in a class of medications called phosphodiesterase (PDE) inhibitors. Sildenafil treats erectile dysfunction by increasing blood flow to the penis during sexual stimulation. This increased blood flow can cause an erection. Sildenafil treats PAH by relaxing the blood vessels in the lungs to allow blood to flow easily.

Product Image Of Sildenafil Citrate 100mg:



How to use sildenafil Citrate 100mg?

Use sildenafil Citrate according to the doctor's instructions. Check whether the label on the drug has an accurate dose description.

Sildenafil is usually injected in your doctor's office, hospital or clinic. If you use sildenafil at home, health care workers will teach you how to use

it. Make sure you know how to use sildenafil. According to the procedure used when the dose is used. If you have any questions, please contact your health care provider.

Do not use sildenafil if it contains particles, turbid or discoloration, or if the vial is broken or damaged.
Regular use of sildenafil can benefit most from it. Use it at the same time every day.
Even if you feel good, continue to use sildenafil. Do not miss any dose.

Do not stop using sildenafil or change your dose suddenly, rather than talk to your doctor.
Keep the product and syringes and needles out of the reach of children and stay away from pets. Do not reuse needles, syringes, or other materials. Please ask your health care staff how to deal with these substances after use. Comply with all local regulations.

If you miss the dose of sildenafil, please use it as soon as possible. If your next dose is almost time, skip the missed dose and return to your regular dosing regimen. Do not use 2 doses at a time.

Does your health care provider ask questions about how to use sildenafil.

Important safety information

Sildenafil may cause drowsiness, dizziness, fainting or blurred vision. If you use alcohol or some drugs, these effects may be worse. Carefully use sildenafil. Do not drive or perform other potentially unsafe tasks until you know your reaction to it.

Sildenafil may cause dizziness, dizziness or fainting; alcohol, hot weather, exercise or fever may increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Sit down or lie down the first sign of any of these effects.

The use of sildenafil in patients with heart disease may increase the risk of heart-related side effects, including heart attacks or strokes. Symptoms of heart attack may include chest, shoulder, neck or jaw pain; arm or leg numbness severe dizziness, headache, nausea, stomach pain or vomiting; fainted or visual changes. Symptoms of stroke may include confusion, vision or speech changes, one-sided weakness or fainting. If you encounter these symptoms, please contact your doctor immediately or seek medical attention immediately.

Sildenafil rarely causes prolonged (more than 4 hours) or painful erections. Even if you do not have sex may also happen. If not treated immediately, may lead to permanent sexual problems, such as yang. The If the erection is more than 4 hours, contact your doctor immediately.

If vomiting or diarrhea occurs, you need to be careful not to dehydrate. Please contact your doctor's instructions.

Sildenafil may not often cause mild, temporary visual changes (eg, blurred vision, sensitivity to light, blue / green tint to vision). If your eyesight is persistent or serious, please contact your doctor.

It is reported that patients taking sildenafil in patients with ocular problems known as non-arterial anterior ischemic optic neuropathy (NAION). In some cases, this may result in loss of vision or permanent loss of vision. If you notice a sudden drop in vision or loss of vision in one eye or both eyes, contact your doctor immediately.

Some patients taking sildenafil have reported a sudden decrease in hearing loss and hearing loss. Sometimes they also notice that the ears are ringing or dizzy.

If you notice sudden reduction or hearing loss, please contact your doctor immediately.

Before accepting any medical or dental care, emergency care or surgery, please tell your doctor or dentist if you take sildenafil.

Do not use drugs or treat ED, while using sildenafil did not first check your doctor.

Carefully use sildenafil in ELDERLY; they may be more sensitive to their impact.
Sildenafil is not approved for use by children. With the increase in sildenafil, the number of child deaths increased. However, your child's doctor may decide that taking the benefits of sildenafil may outweigh the risks. If you have any questions about your child to Sidney, please talk to the doctor.

PREGNANCY and BREAST-FEEDING: If you are pregnant, please contact your doctor. You will need to discuss the benefits and risks of using sildenafil during pregnancy. Do not know if this medicine is found in breast milk.

If you are using sildenafil or are breastfeeding, please consult your doctor. Discuss any risks to your home baby.

Package and Shipping Of Sildenafil Citrate 100mg:

Under 25kg: inner: A PE bag in aluminum foil bag; Outside: carton

Over 25kg: inner: Double PE bags; Outside: 25kg/ fibre drum

Shipping:

With 3-5 days after your payment



FAQ Of Sildenafil Citrate 100mg:

Q1: Do you Accept Sample Order?

A: Yes. The smallest order we accept is 1kg.

Q2: Do you accept VISA business credit card?

A: Yes. VISA business credit card is in our payment.

Q3: Is there any discount?

A: Yes, if you want to purchase larger quantity, we can offer a better price for you.

Q4: How long does it take to me if I order?

A: About 3-5 days upon receipt of payment.

Q5: What kind of transportation do you have?

A: We have many ways of transportations, eg: Air, Sea, Special Lines, TNT, FEDEX, EMS, ect.



Shaanxi Hongbaiyi Biotech Co., Ltd.



18192109180



tracy@sxhongbaiyi.com



peptide-powder.com

Hengjia Business Building, No.115 Weiyang Road, E&T Development Zone, Xi'an, Shaanxi, China.