



Melanotan II Polypeptide Human Growth Hormone Peptide Mt 2 Skin Tanning Peptide 10mg/vial

Our Product Introduction

for more products please visit us on peptide-powder.com

Basic Information

- Place of Origin: China
- Brand Name: Hongbaiyi
- Certification: COA, HPLC MR
- Model Number: HBY-Melanotan 2
- Minimum Order Quantity: 5 box
- Price: US\$ 51-59/box
- Packaging Details: 10mg/vial, 10vials/box
- Delivery Time: 3-5 work days after your payment
- Payment Terms: , Western Union, T/T
- Supply Ability: 3000kg/Month



Product Specification

- Peptide Name: Melanotan 2
- CAS No.: 121062-08-6
- Appearance: White Lyophilized Powder
- Molecular Weight: 1024.2
- Molecular Formula: C₅₀H₆₉N₁₅O₉
- Peptide Purity: 99%min
- Storage: 2-8°C
- Shelf Life: 2 Years



More Images



Product Description

Melanotan II Polypeptide Human Growth Hormone Peptide Mt 2 Skin Tanning Peptide 10mg/vial

Basic Information Form of Melanotan 2

Name	Melanotan 2
CAS NO.	121062-08-6
Purity	99%min
MW	1024.2
MF	C ₅₀ H ₆₉ N ₁₅ O ₉
Appearance	White lyophilized powder
Storage	2-8°C

What Is Melanotan 2

Melanotan is a tanning peptide that stimulates the production of melanin in the body to foster a deep, natural tan. This is the body's way of protecting itself from too much sun exposure by increasing the level of melanin in the body. Melanin is your body's natural response to UV damage. The end result is a darkening of the skin.

Product Image Of Melanotan 2



Application of Melanotan II

Where there is no physical erectile dysfunction and where at least some degree of psychological libido exists, Melanotan II can greatly increase the tendency to penile erection. The situation can even become like the years of puberty, where scarcely even the beginning of a sexual thought or even nothing at all (perhaps an unconscious thought however) produces an erection. General physical fatigue can become irrelevant. These effects will not occur for all individuals, but do occur for many.

Prosexual usage requires injection at least about 4-6 hours prior to desired effect. Duration of action can be about 6-12 hours after onset of effect.

Needed dosage is typically about 1 mg, but for some even 0.5 mg can be effective. As a guide, dosage should not exceed 2 mg.

Because of the associated effect in increasing tendency to tan, frequency of MT-II use for this purpose is generally limited. If wishing this effect frequently, a better choice is to use PT-141 (bremelanotide,) an MT-II derivative.

How is MT2 Used?

This peptide is shipped by vial in a powder dried form that is reconstituted using sterile water and then injected under the skin where it is absorbed into the skin itself and spread throughout.

Administration

Both MT and MT II are currently supplied as white lyophilised powder contained in a sealed multi-use vial. The peptide is

susceptible to temperature degradation and should be shipped preferably with an ice pack though contrary to popular belief, the rate of degradation is very slow (weeks) in its powder form, so there's no need to be alarmed if yours wasn't shipped in this manner or you are unable to collect your package from a depot for a day or two. Once delivered, the powder is best stored in a freezer, or refrigerated if this is not possible.

To prepare for injection, it must be reconstituted with bacteriostatic water. You may use anything between 1ml and 5ml of water for your vial. Dependant upon the amount of water used will determine the concentration of your solution. For example, a 10mg vial of Melanotan II mixed with 1ml of water will provide a solution of 10mg per 1ml (10mg/ml). This means that a 1mg dose will require a shot of 0.1ml. Bearing in mind that the recommended starting dose is 0.25mg, using the example above, the actual volume of the shot would be 0.025ml (1/4 of 1 tenth of a ml). This is a very small volume and very difficult to accurately dose even with a 0.5ml insulin syringe. Therefore, at least until your dosages have increased, it is suggested that you use more water for your vial.

An example of a good solution would be to mix 10mg of Melanotan II powder with 4ml of bacteriostatic water. This now provides: 10mg/4ml or 1mg/0.4ml or 0.25mg/0.1ml. 0.1ml can be accurately measured using a 0.5ml or 1ml syringe.

Package and Shipping of Melanotan 2

Package: 10mg/vial, 10vials/box

Shipping: With 3-5 days after your payment



FAQ of Melanotan 2

Q1: Do you Accept Sample Order?

A: Yes. The smallest order we accept is one box.

Q2: Do you accept VISA business credit card?

A: Yes. VISA business credit card is in our payment.

Q3: Is there any discount?

A: Yes, if you want to purchase larger quantity, we can offer a better price for you.

Q4: How long does it take to me if I order?

A: About 3-5 days upon receipt of payment.

Q5: What kind of transportation do you have?

A: We have many ways of transportations, eg: Air, Sea, Special Lines, TNT, FEDEX, EMS, ect.



Shaanxi Hongbaiyi Biotech Co., Ltd.



18192109180



tracy@sxhongbaiyi.com



peptide-powder.com

Hengjia Business Building, No.115 Weiyang Road, E&T Development Zone, Xi'an, Shaanxi, China.